Physiology of Foveated Human Vision

Retina, Fovea, and Parafovea from Biology to Computation

Luiz Velho, IMPA GPT, Open.AI (collaborator)

Abstract

Human vision is spatially nonuniform. A small central retinal region provides dense cone sampling, low neural convergence, and the highest spatial and chromatic resolution, while progressively more eccentric retina trades detail for sensitivity, coverage, motion processing, and efficiency. This report describes that organization with emphasis on the retina, fovea, foveola, and parafovea. It first follows the transformation from photons to retinal ganglion-cell signals, then explains the anatomical specializations that create the foveal pit and the sharp transition in sampling and circuitry around it. The resulting physiology is connected to visual acuity, crowding, color, temporal sensitivity, active gaze, and engineering models for gaze-contingent sensing and foveated rendering. The main conclusion is that foveation is not simply a high-resolution circular patch. It is a coupled spatial, neural, and motor strategy whose useful engineering analogue must account for eccentricity, content, eye-tracking uncertainty, and time.

Contents

1 Introduction **2**

2 The Eye **2**
2.1 Optical apparatus . . . 2
2.2 Coats, retina, and neural output . . . 3

3 From the retinal image to a neural code **3**
3.1 Image scale at the retina . . . 3
3.2 The retina is neural tissue . . . 4

4 Central retinal geography **4**
4.1 Macula, fovea, foveola, parafovea, and perifovea . . . 4

5 Why the fovea supports high acuity **6**
5.1 The pit clears an optical and neural path . . . 6
5.2 Cone packing and private-line signaling . . . 6

6 The parafovea is a transition regime **6**

7 Foveated vision is active **7**

8 A quantitative language for eccentricity **8**

9 Implications for foveated computation **8**
9.1 What biology supports . . . 8
9.2 Design constraints . . . 8
9.3 Connection to active vision . . . 8

10 Scope and qualifications **9**

11 Conclusions **9**

1 Introduction

The eye is the optical–neural *front end* of the human visual perception system. In a compact engineering description, the complete system comprises (1) the **eye**, which acts as the sensor by focusing light, converting photons into electrical responses, and performing the first stages of neural computation; (2) the **optic nerve and central visual pathways**, which provide the connection from retinal ganglion cells through the optic chiasm, thalamus, and optic radiations; and (3) the **brain**, whose distributed neuronal circuits interpret the signal and connect perception to attention, memory, and action. These are functional divisions rather than isolated modules: the retina is neural tissue, the optic nerve is made of retinal axons, and recurrent brain activity influences what is selected by gaze. The retina is not a uniform camera sensor: its photoreceptor density, interneuron organization, ganglion-cell sampling, vascular structure, and downstream representation all change strongly with retinal eccentricity.

This nonuniform architecture has two consequences. First, the quality of a visual signal depends on where it falls on the retina. The point at which the visual axis meets the central retina is served by the foveal system: a specialized cone mosaic and neural pathway optimized for fine spatial discrimination. Only a few degrees away, the parafovea already exhibits a different balance among acuity, sensitivity, pooling, and contextual processing. Second, stable high-detail perception depends on active gaze. Saccades repeatedly place behaviorally relevant targets on the fovea; fixation, vergence, pursuit, and stabilization mechanisms maintain a useful retinal signal between gaze shifts. The companion report *Eye movements and stereoscopic depth perception* develops those motor and binocular relationships [1]. The present report focuses on the sensor and the central retinal substrate that make foveation valuable.

Vision is one of the dominant computational commitments of the primate cerebrum. In the macaque, more than half of the cerebral cortex is largely or exclusively involved in visual processing, comprising about 32 visual and visual-association areas arranged across approximately ten cortical processing levels [2, 3]. The human system likewise extends far beyond primary visual cortex (V1), through V2, V3/V3A, V4, MT/V5, and higher ventral and dorsal networks. This striking fact is sometimes distorted into the claim that "more than 60% of all brain neurons are dedicated to vision." That literal neuron-count claim is not defensible: of roughly 86 billion neurons in the human brain, about 69 billion are in the cerebellum [4]. The accurate statement concerns the breadth of visual *cortical territory and networks*, not a fixed fraction of every neuron in the brain.

The terms *fovea*, *foveola*, *parafovea*, and *perifovea* have anatomical, clinical, psychophysical, and engineering usages that do not always coincide. Here they are used anatomically unless a broader perceptual region is explicitly stated.

> **Central idea.** Foveated vision is a resource-allocation strategy spanning the entire perception system: a nonuniform ocular sensor, an information-limited optic-nerve connection, extensive cortical processing, and active eye movements.

2 The Eye

The eye is both an optical instrument and a living neural sensor. Its transparent media form an image, its muscular and vascular systems maintain and adapt that image-forming apparatus, and the retina converts light into parallel neural signals. The principal structures and their physiological roles are summarized below and in Figure 1 [5].

2.1 Optical apparatus

The **cornea** is the transparent anterior surface and supplies most of the eye's static refractive power because of the large refractive-index change at the air–tear interface. Behind it, the




Figure 1: Schematic horizontal section of the right eye, viewed from above. The cornea and crystalline lens focus light through the pupil onto the retina; the iris regulates the aperture; the ciliary body changes lens power; and retinal ganglion-cell axons leave at the optic disc to form the optic nerve. Geometry is simplified and not to scale.

aqueous humor fills the anterior and posterior chambers. The **iris** is a contractile diaphragm; its central opening, the **pupil**, sets the light-admitting aperture. Constriction reduces retinal illuminance and many optical aberrations while increasing depth of field, but a very small pupil increases diffraction.

The transparent **crystalline lens** provides the variable component of optical power. The ciliary muscle and zonular fibers change lens curvature during accommodation, allowing near and far objects to be focused on the retina. The large posterior cavity is filled by the transparent vitreous body, which helps maintain globe shape and provides a clear path to the retina. The resulting retinal image is inverted and laterally reversed.

2.2 Coats, retina, and neural output

The tough **sclera** forms the outer structural coat and continues anteriorly as the cornea. Inside it, the vascular, pigmented **choroid** absorbs stray light and supports the metabolically demanding outer retina. The innermost coat is the **retina**: a layered extension of the central nervous system containing photoreceptors, interneurons, ganglion cells, glia, and the retinal pigment epithelium. The macula specializes central vision, and the fovea at its center provides the highest spatial acuity.

The **optic disc** is the exit site for ganglion-cell axons and retinal vessels; because it contains no photoreceptors, it produces the physiological blind spot. The axons form the **optic nerve** and carry retinal output toward the optic chiasm and central targets. The National Eye Institute describes the nerve as a bundle of more than one million fibers [6]. The eye therefore does not send a photograph to the brain: it sends activity in many parallel, feature-selective output channels.

3 From the retinal image to a neural code

3.1 Image scale at the retina

Optical quality is one determinant of resolution, but it is not the only one: photoreceptor spacing, post-receptoral circuitry, fixational motion, and cortical decoding all contribute.

At the posterior pole, approximately 0.28–0.30 mm of retina subtends one degree of visual angle in a typical adult eye. The conversion varies with axial length, so millimeters and degrees should not be treated as universally interchangeable. This matters when mapping histological zones to display coordinates or eye-tracker measurements.

3.2 The retina is neural tissue

The vertebrate retina is inverted: light crosses the inner neural layers before reaching rod and cone outer segments near the retinal pigment epithelium (RPE). Photoreceptors convert absorbed photons into graded electrical responses. In darkness they release glutamate continuously; light hyperpolarizes them and reduces that release. Bipolar cells split the signal into ON and OFF pathways, while horizontal and amacrine cells provide lateral, temporal, and contextual interactions. Ganglion cells are the retinal output neurons; their axons form the optic nerve.




Figure 2: Simplified retinal organization. Light and the principal feed-forward neural signal travel in opposite directions. Lateral circuits are omitted for clarity.

Retinal computation begins before the optic nerve. Center-surround receptive fields emphasize local contrast rather than absolute luminance; parallel ganglion-cell classes differ in spatial scale, temporal response, color opponency, and projection target. The retina therefore transmits a compact, task-relevant description of change and contrast rather than a literal pixel array.

4 Central retinal geography

4.1 Macula, fovea, foveola, parafovea, and perifovea

The macula is the central retinal region associated with high-acuity vision and macular pigment. Within it, the fovea is a shallow pit centered on the visual axis. The foveola is the central floor of that pit. The parafovea surrounds the fovea and contains the greatest accumulation of displaced inner retinal neurons; the perifovea is the next outward annulus. Common approximate dimensions are summarized in Table 1. They are conventions, not exact biological boundaries, and natural inter-individual variation is substantial [7–9].

In psychophysics and computer graphics, *parafoveal* is often used more loosely for a few degrees around fixation, and *peripheral* may begin wherever a particular task shows a substantial decline. Any engineering specification should therefore state the eccentricity range in degrees rather than rely only on region names.

Table 1: Approximate anatomical scales near the human visual axis. Angular values assume about 0.29 mm per degree and are rounded.

Region	Approximate linear extent	Approximate visual extent	Functional emphasis
Foveola	central 0.35 mm diameter	about 1.2 degrees	Densest cone sampling; no rods at the center; highest spatial resolution
Fovea	about 1.5 mm diameter	about 5 degrees	Pit and sloping walls; inner layers displaced; cone-dominated central vision
Parafovea	about 0.5 mm-wide annulus	roughly 5–8 degrees diameter	Thick inner retina; falling cone density; rising rod contribution and neural pooling
Perifovea	next roughly 1.5 mm annulus	out to roughly 18 degrees diameter	Transition toward peripheral retinal organization
Macula	about 5–6 mm diameter	roughly central 18–20 degrees	Broad central retina containing all zones above




Figure 3: Nested central retinal zones shown as a top-down schematic. The areas are intentionally not drawn to the same scale as the labels.

5 Why the fovea supports high acuity

5.1 The pit clears an optical and neural path

During development, ganglion cells, bipolar cells, and their processes are displaced laterally from the center, while cone photoreceptors migrate inward and elongate. The mature pit therefore leaves a comparatively direct optical path to the central cones and creates the radially oriented Henle fiber layer that connects cone pedicles to laterally displaced bipolar cells [7, 8]. The central foveal avascular zone removes capillaries from the optical path, although the tissue still requires intense metabolic support from the choroidal circulation and surrounding retinal capillary networks.

The pit itself is not sufficient for high acuity. People with foveal hypoplasia can still show central cone elongation and packing, and foveal morphology varies across healthy observers. High-resolution vision emerges from several aligned adaptations: optical quality, narrow and elongated cone apertures, dense packing, low convergence in midget pathways, stable fixation, and expanded cortical representation.

5.2 Cone packing and private-line signaling

Human cone density reaches a sharp and individually variable maximum near the foveal center. Histological measurements reported peak values approaching 200,000 cones/mm^2, followed by a steep decline with eccentricity [10]. At the very center, rod photoreceptors are absent; rods appear outside the rod-free zone and become increasingly important with eccentricity. The distribution is therefore not a simple trade between "good" and "poor" vision. Central cones favor fine spatial and chromatic sampling under photopic conditions, while rods and larger receptive fields increase sensitivity and support dim-light detection away from the center.

In central midget circuitry, a single cone can feed a single midget bipolar cell and then a single midget ganglion cell, especially near the fovea [7, 11]. This low convergence preserves the spatial identity of the input. With increasing eccentricity, more photoreceptors contribute to downstream receptive fields, improving sensitivity and efficiency while reducing spatial specificity.




Figure 4: Conceptual change in convergence with eccentricity. Real retinal circuits contain multiple parallel cell classes and are more heterogeneous than this schematic.

6 The parafovea is a transition regime

The parafovea should not be described merely as a blurred version of the fovea. Anatomically, it contains the lateral displacement generated by foveal construction: inner nuclear and ganglion-cell layers thicken on the foveal slope and rim. Functionally, it combines still-substantial cone vision with rapidly changing cone spacing, increasing rod contribution, larger receptive fields, more convergence, and stronger effects of surrounding context.

This transition changes several perceptual dimensions at different rates:

- **Spatial resolution.** Minimum resolvable detail increases with eccentricity. Over useful ranges, many size thresholds can be approximated by a linear function of eccentricity after a task-specific intercept [12, 13].
- **Crowding.** Identification of an object can fail because nearby flankers interfere, even when the isolated object remains detectable. Critical spacing grows approximately with eccentricity, so peripheral recognition cannot be modeled by blur alone [13, 14].
- **Color and form.** Fine chromatic discrimination is strongest centrally, but color is not absent in the periphery. Performance depends on stimulus size, luminance, adaptation, and retinal location.
- **Sensitivity and temporal processing.** Larger pooling areas and rod-rich retina can favor detection under low light and support wide-field motion and event sensitivity, even as fine spatial detail declines.
- **Binocular function.** Stereoacuity is best near fixation. Eccentricity, spatial frequency, contrast, crowding, and eye alignment all affect the usable disparity signal.




Figure 5: Conceptual profiles with eccentricity. The curves illustrate different trends, not a shared physiological unit or fitted dataset. Exact functions depend on meridian, observer, adaptation, and task.

7 Foveated vision is active

The useful field of view is far larger than the high-acuity foveal field. The oculomotor system resolves this mismatch by moving the eyes. Peripheral and parafoveal signals provide target candidates through contrast, motion, novelty, task relevance, and coarse scene structure. A saccade relocates the foveae; fixation allows high-resolution analysis; pursuit and vestibulo-ocular responses stabilize selected content; and vergence aligns the two eyes in depth. Fixational drift and microsaccades continue to transform the retinal input even during attempted steady gaze [15].

This recurrent organization can be written as a perception-action cycle:

peripheral evidence → target selection → gaze shift → foveal analysis → updated task state.

The scene is not necessarily stored as a uniformly detailed internal image. Stable perception can arise from task-dependent representations, memory, prediction, and the availability of detail through another gaze shift. The biological strategy is thus closer to an active query system than to a fixed megapixel camera.

8 A quantitative language for eccentricity

Let e denote angular eccentricity from the current fixation. A common psychophysical approximation expresses a threshold size $s(e)$ as

$$s(e) \approx s_0 \left(1 + \frac{e}{E_2}\right), \tag{1}$$

where s_0 is the foveal threshold and E_2 is the eccentricity at which the threshold doubles. The parameter is not a universal constant: it changes with task, contrast, orientation, temporal conditions, and meridian. Equation 1 is therefore a coordinate transform for a specific measured capability, not a full model of peripheral vision.

Retinal ganglion-cell receptive-field density can also be modeled as a function of visual-field position [16]. Such models improve on circular three-zone approximations because they can encode meridional asymmetry and continuous falloff. Even then, retinal sampling alone does not predict perception. Optical blur, fixational instability, cortical magnification, attention, and crowding remain important.

9 Implications for foveated computation

9.1 What biology supports

Foveated rendering and sensing allocate computation according to gaze. The biological motivation is sound: spatial detail requirements decline rapidly away from fixation, and wide-field systems otherwise spend large resources on samples the observer cannot resolve. Early multiresolution systems and modern gaze-contingent graphics exploit this principle [17–19].

The closest engineering abstraction is a continuous, anisotropic, time-dependent sampling field centered on estimated binocular gaze. A practical implementation may still use concentric regions, but their boundaries should be understood as approximations to several overlapping functions rather than literal retinal anatomy.

9.2 Design constraints

If the predicted gaze at display time is $\hat{g}(t + \Delta)$ and the combined angular uncertainty is σ_g, a conservative foveal radius can be expressed schematically as

$$r_{\text{HQ}} = r_{\text{task}} + k\sigma_g + v_{\text{eye}}\Delta, \tag{2}$$

where r_{task} is the radius needed by the visual task, v_{eye} represents plausible gaze velocity during the prediction interval, and k sets the desired confidence margin. This is not a physiological law; it is a useful engineering decomposition of content demand, uncertainty, and latency.

9.3 Connection to active vision

The broader lesson for active vision is to optimize the next observation, not merely compress the current one. A biological or artificial observer can direct its high-resolution sensor toward locations that reduce uncertainty, disambiguate depth, verify an object hypothesis, or support an action. In that sense, foveation is both an encoding strategy and a control policy.

Table 2: Biological facts and their engineering consequences.

Biological fact	Engineering interpretation	Common failure mode
Resolution falls continuously with eccentricity	Use smooth, task-calibrated quality falloff	Visible rings or abrupt changes during small gaze errors
Different capabilities fall off differently	Separate models for acuity, contrast, color, motion, and disparity	Treating peripheral vision as uniform Gaussian blur
Crowding limits recognition	Preserve feature spacing, contrast structure, and semantic salience	Readable isolated details become unrecognizable in clutter
Gaze changes rapidly during saccades	Predict gaze and manage temporal transitions	Post-saccadic exposure of a low-quality region
Eye tracking is noisy and delayed	Expand the high-quality region by an uncertainty and latency margin	Quality boundary enters conscious view
The two eyes may differ in direction and depth	Estimate binocular gaze and vergence, not only a cyclopean 2-D point	Interocular mismatch and unstable stereo detail
Peripheral vision remains useful	Preserve motion, transients, contrast, and scene layout	Tunnel vision despite acceptable spatial-resolution metrics

10 Scope and qualifications

Several simplifications should remain explicit.

- Histological boundaries do not create step changes in perception.
- Peak cone density, foveal pit shape, avascular-zone size, axial length, and visual performance vary across observers.
- Acuity is not a single number: detection, resolution, recognition, vernier alignment, stereoacuity, and reading have different eccentricity functions.
- Peripheral vision is not defined only by reduced spatial frequency. Crowding, uncertainty, pooling, attention, and temporal sensitivity matter.
- A gaze tracker measures an estimate of eye orientation, not direct access to the observer's attended object or exact retinal locus.
- Three-ring rendering profiles are implementation conveniences, not faithful anatomical models of fovea, parafovea, and periphery.

11 Conclusions

The retina is a layered neural processor whose sampling strategy changes radically with eccentricity. The foveal center combines dense, elongated cones, displaced inner layers, an avascular optical path, low-convergence midget circuitry, and expanded downstream representation. The parafovea is the steep transition around that specialization: cone density falls, rods and pooling increase, inner layers thicken, and perception becomes progressively more dependent on context and crowding. Peripheral retina then extends coverage and sensitivity with coarser spatial sampling.

Eye movements turn this nonuniform sensor into an effective visual system. Peripheral evidence selects targets, saccades relocate the fovea, stabilization preserves useful signals, and successive fixations support task-dependent understanding. Foveated computation should preserve this logic: allocate resources continuously with eccentricity, protect temporal transitions, account for tracking uncertainty and binocular state, and retain the peripheral signals that guide the next gaze shift.

Source note

This report was developed as a companion to the document *Eye movements and stereoscopic depth perception* [1]. Its treatment of saccades, fixation, vergence, and active depth inference motivated the perception-action framing. The anatomical and physiological material was expanded using the literature cited below.

References

[1] Eye movements and stereoscopic depth perception, 2026.

[2] James J. Knierim and David C. Van Essen. Visual cortex: cartography, connectivity, and concurrent processing. *Current Opinion in Neurobiology*, 2(2):150–155, 1992. doi: 10.1016/0959-4388(92)90003-4.

[3] Daniel J. Felleman and David C. Van Essen. Distributed hierarchical processing in the primate cerebral cortex. *Cerebral Cortex*, 1(1):1–47, 1991. doi: 10.1093/cercor/1.1.1-a.

[4] Frederico A. C. Azevedo, Ludmila R. B. Carvalho, Lea T. Grinberg, José M. Farfel, Renata E. L. Ferretti, Renata E. P. Leite, Wilson Jacob Filho, Roberto Lent, and Suzana Herculano-Houzel. Equal numbers of neuronal and nonneuronal cells make the human brain an isometrically scaled-up primate brain. *Journal of Comparative Neurology*, 513(5):532–541, 2009. doi: 10.1002/cne.21974.

[5] National Eye Institute. How the eyes work. `https://www.nei.nih.gov/eye-health-information/healthy-vision/how-eyes-work`, 2022. Last reviewed April 20, 2022; accessed September 3, 2026.

[6] National Eye Institute. About the eye. `https://www.nei.nih.gov/eye-health-information/healthy-vision/nei-for-kids/about-eye`, 2025. Last reviewed December 10, 2025; accessed September 3, 2026.

[7] Jan M. Provis, Adam M. Dubis, Ted Maddess, and Joseph Carroll. Adaptation of the central retina for high acuity vision: cones, the fovea and the avascular zone. *Progress in Retinal and Eye Research*, 35:63–81, 2013. doi: 10.1016/j.preteyeres.2013.01.005.

[8] Andreas Bringmann, Steffen Syrbe, Katja Gorner, Johannes Kacza, Mike Francke, Peter Wiedemann, and Andreas Reichenbach. The primate fovea: Structure, function and development. *Progress in Retinal and Eye Research*, 66:49–84, 2018. doi: 10.1016/j.preteyeres.2018.03.006.

[9] Adam M. Dubis, Dean M. Costakos, Charu D. Subramaniam, Pooja Godara, William J. Wirostko, Joseph Carroll, and Jan M. Provis. Evaluation of normal human foveal development using optical coherence tomography and histologic examination. *Archives of Ophthalmology*, 130(10):1291–1300, 2012. doi: 10.1001/archophthalmol.2012.2270.

[10] Christine A. Curcio, Kenneth R. Sloan, Robert E. Kalina, and Anita E. Hendrickson. Human photoreceptor topography. *Journal of Comparative Neurology*, 292(4):497–523, 1990. doi: 10.1002/cne.902920402.

[11] Dennis M. Dacey. The mosaic of midget ganglion cells in the human retina. *Journal of Neuroscience*, 13(12):5334–5355, 1993. doi: 10.1523/JNEUROSCI.13-12-05334.1993.

[12] Stuart M. Anstis. A chart demonstrating variations in acuity with retinal position. *Vision Research*, 14(7):589–592, 1974. doi: 10.1016/0042-6989(74)90049-2.

[13] Hans Strasburger, Ingo Rentschler, and Martin Juttner. Peripheral vision and pattern recognition: A review. *Journal of Vision*, 11(5):13, 2011. doi: 10.1167/11.5.13.

[14] Ruth Rosenholtz. Capabilities and limitations of peripheral vision. *Annual Review of Vision Science*, 2:437–457, 2016. doi: 10.1146/annurev-vision-082114-035733.

[15] Michele Rucci and Jonathan D. Victor. The unsteady eye: an information-processing stage, not a bug. *Trends in Neurosciences*, 38(4):195–206, 2015. doi: 10.1016/j.tins.2015.01.005.

[16] Andrew B. Watson. A formula for human retinal ganglion cell receptive field density as a function of visual field location. *Journal of Vision*, 14(7):15, 2014. doi: 10.1167/14.7.15.

[17] Wilson S. Geisler and Jeffrey S. Perry. Real-time foveated multiresolution system for low-bandwidth video communication. In *Human Vision and Electronic Imaging III*, volume 3299, pages 294–305. SPIE, 1998. doi: 10.1117/12.320120.

[18] Brian Guenter, Mark Finch, Steven Drucker, Desney Tan, and John Snyder. Foveated 3d graphics. *ACM Transactions on Graphics*, 31(6):164:1–164:10, 2012. doi: 10.1145/2366145.2366183.

[19] Anjul Patney, Marco Salvi, Joohwan Kim, Anton Kaplanyan, Chris Wyman, Nir Benty, David Luebke, and Aaron Lefohn. Towards foveated rendering for gaze-tracked virtual reality. *ACM Transactions on Graphics*, 35(6):179:1–179:12, 2016. doi: 10.1145/2980179.2980246.